

# Fermions in Deep Optical Lattice under p-wave Feshbach Resonance

S.-K. Yip

*Institute of Physics, Academia Sinica, Nankang, Taipei 115, Taiwan*

(Dated: February 2, 2008)

## Abstract

We investigate theoretically the energy levels of two identical Fermions in a harmonic potential well under p-wave Feshbach resonance. The magnetic fields needed to affect the energy levels differ strongly among the levels, and they can be at values far from the free space resonant field.

PACS numbers: 03.75.-b, 71.10.Ca, 37.10.Jk

arXiv:0801.0636v1 [cond-mat.other] 4 Jan 2008

There are substantial recent activities in studying cold atomic gases in the presence of an optical lattice (e.g. [1]). Part of this interest stems from the possible connection with solid-state physics (e.g. [2]) and the vast theoretical literature on exotic states in lattice hamiltonians. Generally, the interaction between atoms are weak. To increase the interaction between particles, one possible way is to make use of Feshbach resonance. Indeed, a few such experiments have already been carried out [3, 4]. An important question arises already even when the tunneling of atoms between different potential wells can be neglected. For more than one particle in a well, the energy levels, wavefunctions etc are strongly modified by the Feshbach resonance. Theoretical studies of this "dressed state" have been carried out in some recent papers [5, 6]. Experimentally, Ref [3] has been able to obtain some important information about this dressed state, namely, the probability of finding a particle in a particular harmonic oscillator level. They did this by starting with a deep lattice potential and, by sweeping the magnetic field through resonance and then turning off the optical lattice, occupation of the different Bloch bands was experimentally observed. The energy of the lowest bound state has also been measured by radio frequency spectroscopy [4]. Understanding these dressed state is also crucial for obtaining the lattice Hamiltonian [7], the first essential step to eventually able to understand any many-body systems to be studied.

The above references [3, 4, 5, 6, 7] are on s-wave Feshbach resonance between two Fermions of different internal hyperfine spins. Here we would like to extend the study to two identical fermions (same hyperfine spin) under a p-wave Feshbach resonance. P-wave Feshbach resonances in Fermions (e.g. [8, 9, 10]) itself have generated much activities. One reason is possible p-wave superfluidity (e.g. [11, 12, 13, 14]). There has also been a suggestion [15] to obtain molecular superfluid by first starting with a lattice with two atoms per site and then "melt" the Mott state by turning off the optical lattice. This may perhaps be a route to obtain p-wave superfluidity and gives additional impetus to study Fermionic atoms under p-wave Feshbach resonance within an optical lattice.

In this paper, we shall then consider two identical Fermions in a harmonic potential well under a p-wave Feshbach resonance. We shall find how the energy levels depend on the magnetic field detuning. We shall show that, in contrast to s-wave resonances, due to the strong momentum dependence of the effective interaction, there is a large shift in the magnetic field from resonance which is necessary to affect the energy levels with the shift

itself depending strongly on the energy level under study. The validity of this prediction can be tested by performing an experiment similar to that of [3]. We employed a method different from the existing literature, an approach some readers may find more transparent. We shall also report on the generalization to resonance with higher angular momenta.

In a harmonic trap, we can first separate the center of mass versus the relative motion, and we need only to treat the latter. The wavefunction of the "dressed" molecule in the relative coordinate  $\vec{r}$  consists of two parts,  $\psi(\vec{r})$  in the open channel and  $\gamma\varphi_c(\vec{r})$  in the closed channel. Here, for later convenience, we have introduced a normalized wavefunction  $\varphi_c(\vec{r})$  in the closed channel and denote the amplitude of our dressed molecule in this channel by  $\gamma$ . The normalization of the full wavefunction is thus  $|\gamma|^2 + \int_{\vec{r}} |\psi(\vec{r})|^2 = 1$ , where  $\int_{\vec{r}}$  is the short-hand for  $\int d^3\vec{r}$ . For p-wave resonance, due to the dipole interaction, the resonant field depends on the angular momentum  $m_l$  along the applied magnetic field direction. For example, for  $^{40}\text{K}$  in the  $|\frac{9}{2}, -\frac{7}{2}\rangle$  hyperfine state, the  $m_l = 0$  resonance occurs at a field around 0.5G above the  $m_l = \pm 1$  ones [8]. For definiteness, in below we shall assume that we have the  $m_l = 0$  resonance only. Modifications need to other  $m_l$ 's are evident.

The Schrödinger equations are

$$\left(-\frac{\hbar^2}{2m_r}\nabla^2 + U(\vec{r})\right)\psi(\vec{r}) + W(\vec{r})\gamma\varphi_c(\vec{r}) = E\psi(\vec{r}) \quad (1)$$

$$W(\vec{r})\psi(\vec{r}) + \nu'\gamma\varphi_c(\vec{r}) = E\gamma\varphi_c(\vec{r}) \quad (2)$$

Here  $m_r = m/2$  is the reduced mass,  $U(\vec{r})$  is the trap potential, and  $\varphi_c(\vec{r})$  is the closed-channel molecule of  $l = 1$ ,  $m_l = 0$  symmetry with (unrenormalized) energy  $\nu'$ , and  $W(\vec{r})$  is the coupling between the open and closed channels, and  $E$  is the energy of the dressed state. Here, we have already truncated the Hilbert space for the closed-channel to one single state responsible for the resonance of interest. For simplicity, we have also suppressed the part of the wavefunction representing the internal (hyperfine) degrees of freedom which distinguishes between the open and closed channels. Eq (1-2) are the same as those written down for the s-wave (e.g. [5]) except the symmetry of  $\varphi_c(\vec{r})$ .

Equation (1) can be solved in terms of the Green's function defined by

$$\left(-\frac{\hbar^2}{2m_r}\nabla^2 + U(\vec{r}) - E\right)G_E(\vec{r}, \vec{r}') = \frac{2\pi}{m_r}\delta^3(\vec{r} - \vec{r}') . \quad (3)$$

Substituting the result into eq (2), we obtain the implicit equation for the energy  $E$  as

$$E - \nu' + \frac{2\pi}{m_r} \int_{\vec{r}, \vec{r}'} \tilde{\varphi}_c^*(\vec{r}) G_E(\vec{r}, \vec{r}') \tilde{\varphi}_c(\vec{r}') = 0 \quad (4)$$

where we have introduced the short-hand  $\tilde{\varphi}_c(\vec{r}) \equiv W(\vec{r})\varphi_c(\vec{r})$ . The problem thus reduces to finding  $G_E(\vec{r}, \vec{r}')$ . We here adopt a different method from ref [16]. We would also like to express our final answer in terms of experimentally available quantities. For the latter, we first observe that, in the absence of the trapping potential, the expression in eq (4) is related to a quantity that appears in the scattering amplitude. Explicitly, the scattering amplitude for incoming (outgoing) wavevector  $\vec{k}$  ( $\vec{k}'$ ) can be found to be

$$f_{\vec{k}}(\vec{k}') = \frac{-\frac{2\pi}{m_r}\tilde{\varphi}_c^*(\vec{k})\tilde{\varphi}_c(\vec{k}')}{E - \nu' + \frac{2\pi}{m_r}\int_{\vec{r}, \vec{r}'}\tilde{\varphi}_c^*(\vec{r})G_E^{free}(\vec{r}, \vec{r}')\tilde{\varphi}_c(\vec{r}')} \quad (5)$$

where  $G_E^{free}(\vec{r}, \vec{r}')$  obeys eq (3) with  $U(\vec{r})$  set to zero. Here  $E \equiv \frac{\hbar^2 k^2}{2m_r}$ ,  $\tilde{\varphi}_c(\vec{k}) \equiv \int_{\vec{r}}\tilde{\varphi}_c(\vec{r})e^{-i\vec{k}\cdot\vec{r}}$ . For scattering in the  $l = 1$ ,  $m_l = 0$  channel,  $f_{\vec{k}}(\vec{k}') = 4\pi f_{10}(k)Y_1^{0*}(\hat{k})Y_1^0(\hat{k}')$  where  $f_{10}$  has the following form at small energies:

$$f_{10}(k) = \frac{k^2}{-\frac{1}{v} + ck^2 - ik^3}. \quad (6)$$

Here  $v$  has the dimension of a volume and  $c$  inverse length. These parameters, in principle available from experiment (e.g. [8]) or other theoretical calculations, will be used as an input below. Near the resonant field  $B_0^*$ ,  $-\frac{1}{v}$  is roughly linear in  $(B - B_0^*)$ , whereas  $c$  can be regarded as a constant.

At small energies,  $\tilde{\varphi}_c(\vec{k}) \approx -ik_z \int_{\vec{r}}\tilde{\varphi}_c(\vec{r})z \equiv -ik_z\sqrt{3}\alpha$ , where we have also defined the coupling constant  $\alpha$  (This definition is identical with that in [13, 14]). We then have the correspondence

$$-\frac{m_r}{2\pi|\alpha|^2}\left[E - \nu' + \frac{2\pi}{m_r}\int_{\vec{r}, \vec{r}'}\tilde{\varphi}_c^*(\vec{r})G_E^{free}(\vec{r}, \vec{r}')\tilde{\varphi}_c(\vec{r}')\right] = \left[-\frac{1}{v} + ck^2 - ik^3\right] \quad (7)$$

This equation, when expanded and comparing powers of  $k$ , yields the same "renormalization" relations as those written before in momentum space in, e.g., [11, 13, 14].

The Green's function  $G_E(\vec{r}, \vec{r}')$  and  $G_E^{free}(\vec{r}, \vec{r}')$  are both singular as  $\vec{r} \rightarrow \vec{r}'$ , diverging as  $\frac{1}{|\vec{r}-\vec{r}'|}$ , but otherwise finite. Defining  $G_E^{reg}(\vec{r}, \vec{r}') \equiv G_E(\vec{r}, \vec{r}') - G_E^{free}(\vec{r}, \vec{r}')$ ,  $G_E^{reg}(\vec{r}, \vec{r}')$  is thus finite for all  $\vec{r}$  and  $\vec{r}'$ . The equation (4) for the energy can then be written as

$$-\frac{1}{v} + ck^2 - ik^3 = \frac{1}{|\alpha|^2}\int_{\vec{r}, \vec{r}'}\tilde{\varphi}_c^*(\vec{r})G_E^{reg}(\vec{r}, \vec{r}')\tilde{\varphi}_c(\vec{r}') \quad (8)$$

It can be verified that  $G_E^{reg}$  obeys

$$\left(-\frac{\hbar^2}{2m_r}\nabla^2 + U(\vec{r}) - E\right)G_E^{reg}(\vec{r}, \vec{r}') = -U(\vec{r})G_E^{free}(\vec{r}, \vec{r}'). \quad (9)$$

Below, we shall find  $G_E^{reg}(\vec{r}, \vec{r}')$  for the case of a spherical trap of frequency  $\omega$ , thus  $U(\vec{r}) = \frac{1}{2}m_r\omega^2r^2$ . It is sufficient to do this for  $r > r'$  since the  $r < r'$  part can be obtained by the symmetry of  $G_E^{reg}(\vec{r}, \vec{r}')$ . We have  $G_E^{free}(\vec{r}, \vec{r}') = \frac{e^{ik|\vec{r}-\vec{r}'|}}{|\vec{r}-\vec{r}'|} = 4\pi ik \sum_{l, m_l} j_l(kr')h_l^{(1)}(kr)Y_{lm_l}(\hat{r})Y_{lm_l}^*(\hat{r}')$  for  $r > r'$ . By the symmetry of  $\tilde{\varphi}_c$ , we see that only the terms with  $l = 1$ ,  $m_l = 0$  would contribute. Writing this part of  $G_E^{reg}(\vec{r}, \vec{r}')$  as  $g_1(r, r')Y_{10}(\hat{r})j_1(kr')Y_{10}^*(\hat{r}')$ , we are thus left to solve

$$\left( -\frac{\hbar^2}{2m_r} \frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} + \frac{2}{2m_r r^2} + U(r) - E \right) g_1(r, r') = -U(r)4\pi ik h_1^{(1)}(kr) \quad (10)$$

Fortunately, a particular solution for eq (10) can easily be seen to be  $-4\pi ik h_1^{(1)}(kr)$ . The origin of this is simply that  $-G_E^{free}(\vec{r}, \vec{r}')$  is a particular solution to eq (9). The homogeneous solution to eq (10) can be expressed in terms of the confluent hypergeometric functions  $F$ , so we obtain thus

$$g_1(r, r') = \left[ b_1(r') \frac{r}{l_r} F(-\mu_1, \frac{5}{2}, \left(\frac{r}{l_r}\right)^2) + b_2(r') \left(\frac{l_r}{r}\right)^2 F(-\mu_2, -\frac{1}{2}, \left(\frac{r}{l_r}\right)^2) \right] e^{-\frac{1}{2}\left(\frac{r}{l_r}\right)^2} - 4\pi ik h_1^{(1)}(kr) \quad (11)$$

where  $\mu_1 \equiv \frac{E}{2\omega} - \frac{5}{4}$ ,  $\mu_2 \equiv \frac{E}{2\omega} + \frac{1}{4} = \mu_1 + \frac{3}{2}$ , and  $l_r \equiv \sqrt{\hbar/m_r\omega}$  is the oscillator length for relative motion, and  $b_1(r')$  and  $b_2(r')$  are coefficients to be determined. The condition that  $G_E(\vec{r}, \vec{r}') \rightarrow 0$  as  $r \rightarrow \infty$  gives  $b_1 \frac{\Gamma(5/2)}{\Gamma(-\mu_1)} + b_2 \frac{\Gamma(-1/2)}{\Gamma(-\mu_2)} = 0$ , that is  $b_1 = \frac{8}{3} \frac{\Gamma(-\mu_1)}{\Gamma(-\mu_2)} b_2$ . For small  $r'$ ,  $j_1(kr') \rightarrow kr'/3$ . The condition that  $G_E(\vec{r}, \vec{r}')$  must be regular in the limit  $\lim_{r' \rightarrow 0} \lim_{r \rightarrow r'_+}$  therefore requires that there cannot be any  $1/r^2$  term in  $g_1(r, r')$ . Using  $h_1^{(1)}(kr) \rightarrow -i/(kr)^2$  as  $r \rightarrow 0$ , we therefore have  $\lim_{r' \rightarrow 0} b_2(r') = \frac{4\pi}{kl_r^2}$ . On the other hand, for small  $r$ , we obtain from eq (11) that  $g_1(r, r') = b_1(r') \frac{r}{l_r} - \frac{4\pi}{3} ik^2 r$ . (It can be verified that, with the above value of  $b_2$ , there are no terms proportional to  $r^{-1}$  or  $r^0$  in  $g_1(r, r')$ ). Substituting this expression into the right-hand-side (RHS) of eq (8), we see that, in the limit of short-ranged  $\tilde{\varphi}_c$ , we need only the value of  $b_1(r' \rightarrow 0)$ , and the RHS can be evaluated to be  $\frac{3k}{4\pi} \frac{b_1(0)}{l_r} - ik^3$ . The  $-ik^3$  terms on the two sides of eq (8) cancel. Using the above obtained value for  $b_1(0)$  and the definition for  $\mu_1$  and  $\mu_2$ , we obtain finally the equation

$$-\frac{1}{v} + ck^2 = \frac{8}{l_r^3} \frac{\Gamma(\frac{5}{4} - \frac{E}{2\omega})}{\Gamma(-\frac{1}{4} - \frac{E}{2\omega})} \quad (12)$$

for the energies of the dressed states, expressed entirely in terms of quantities entering the scattering amplitude (6) and the harmonic trap frequency  $\omega$  ( $k^2 = 2m_r E$ ).

It is convenient to rewrite this equation in dimensionless form:

$$-\frac{l_r^3}{v} - (-2cl_r)\frac{E}{\omega} = 8\frac{\Gamma(\frac{5}{4} - \frac{E}{2\omega})}{\Gamma(-\frac{1}{4} - \frac{E}{2\omega})} \quad (13)$$

For [8]  $^{40}\text{K}$   $|\frac{9}{2}, -\frac{7}{2}\rangle$ ,  $-c \approx 0.02a_0^{-1}$ , whereas  $l_r \approx 1600a_0$  for a trap of frequency  $\omega = 70\text{kHz}$ , a typical experimental number. Thus the product  $(-2cl_r)$  is large and  $\approx 64$ . The resulting energy levels are plotted in Fig 1. For "high" magnetic field,  $(-\frac{1}{v} \rightarrow \infty)$ , the energy levels are given by the non-interacting values  $(2n + \frac{5}{2})\omega$  ( $n$  is a non-negative integer). For  $n = 0$ , the state has energy decreasing with field, becomes linear in  $-1/v$  for  $-1/v \lesssim 100$ , and thus becomes indistinguishable with the the free space bound state energy [11, 13, 14]  $-\epsilon_b = -\frac{\hbar^2}{2m_r(-cv)}$  for  $B < B_0^*$ . For  $n \geq 1$  and decreasing field, the energy of the level shift to  $(2n + \frac{1}{2})\omega = (2(n-1) + \frac{5}{2})\omega$ . However, the value of the field where the energy crosses from its high field value to the low field one depends on the level  $n$ , increasing as  $n$  increases. The width of field for this transition also increases with  $n$ . This is due to the large  $(-cl_r)$  value discussed above.

The energy levels can be measured by radio-frequency spectroscopy [4]. Below, we analyze instead the implication on an experiment similar to that in [3] (theoretically analyzed in [6, 17]), summarized already briefly in the introduction. We thus evaluate, for a given energy level of Fig 1, the probability of finding a particle in the single particle harmonic oscillator states. For this, we first have to express the two-particle wavefunction  $\psi(\vec{r})$  in harmonic oscillator basis. A procedure similar to that in [6] gives  $\psi(\vec{r}) = \sum_n \eta_n \psi_{n10}(\vec{r})$ , where

$$\eta_n = \frac{A_n / (\frac{E}{2\omega} - (2n + \frac{5}{2}))}{\left\{ \frac{\pi^{3/2} l_r^5}{3|\alpha|^2} + \sum_n [A_n / (\frac{E}{2\omega} - (2n + \frac{5}{2}))]^2 \right\}^{1/2}}, \quad (14)$$

$\psi_{n10}(\vec{r}) \equiv \frac{A_n}{\pi^{3/4} l_r^{5/2}} r \cos\theta F(-n, \frac{5}{2}, (\frac{r}{l_r})^2) e^{-\frac{1}{2}(\frac{r}{l_r})^2}$  are the normalized wavefunctions with  $l = 1$ ,  $m_l = 0$ ,  $A_n \equiv \left[ \frac{1}{3} \frac{(2n+3)!}{2^{n-1} n!} \right]^{1/2}$  are normalization coefficients. The amplitude in the closed channel is  $\gamma = [1 - \sum_n |\eta_n|^2]^{1/2}$ . Below, we consider the special case where the term  $\propto \frac{1}{\alpha^2}$  in the denominator of eq (14) is negligible, correspondingly  $\gamma \rightarrow 0$ . (This assumption affects only quantitatively the probabilities at intermediate fields given below but not the qualitative conclusions.) Using this  $\psi(\vec{r})$  and assuming that the center of mass motion is in its harmonic oscillator ground state, it is straight-forward to express the resulting wavefunction in the coordinates  $\vec{r}_1, \vec{r}_2$  of the two particles. For the latter wavefunction  $\phi$ , we shall employ

instead the quantum numbers  $n_x$ ,  $n_y$  and  $n_z$  since these are more directly related to the occupation of the different Bloch bands and thus the momentum states after turning off of the optical lattice. For example, the state with quantum numbers  $n_x$ ,  $n_y$ ,  $n_z$  contributes to momentum states with  $n_{x(y,z)} \frac{\pi}{d} < |k_{x(y,z)}| < (n_{x(y,z)} + 1) \frac{\pi}{d}$ , where  $d$  is the distance between neighboring sites in the lattice [6]. The resulting expressions however are lengthy and we shall not write them down here, but only give the probabilities below. For definiteness, we consider an initial field  $B \ll B_0^*$  with initial two-particle state corresponding to relative motion with quantum numbers  $n = 0, l = 1, m = 0$  (Energy of relative motion =  $\frac{5}{2}\omega$ .) In the two-particle basis, the state is given by  $\frac{1}{\sqrt{2}} [\phi_{000}(\vec{r}_1)\phi_{001}(\vec{r}_2) - \phi_{000}(\vec{r}_2)\phi_{001}(\vec{r}_1)]$  where the subscripts stands for  $(n_x, n_y, n_z) \equiv \vec{n}$ . The initial probabilities of finding a particle in the single-particle state  $\phi_{\vec{n}}$  are given by  $P_{000} = 1, P_{001} = 1$ . ( $P_{\vec{n}} = 0$  for all other  $\vec{n}$ ). The final probabilities of finding a particle in state  $\vec{n}$  when the magnetic field is swept to the final value  $B$  where the corresponding scattering parameter is  $1/v$  is plotted for a few  $\vec{n}$ 's in Fig 2. The transition from the "low" field values to the "high" field values occur around  $-l_r^3/v \approx 200$  (*c.f.* Fig 1) for this particular state (This field increases with  $n$ ). For magnetic fields with  $-1/v$  above this value, the probabilities correspond to the two-particle relative motion in the  $n = 1, l = 1, m = 0$  state. We find  $P_{000} = \frac{1}{4}, P_{001} = \frac{11}{20}, P_{002} = \frac{9}{20}, P_{100} = P_{010} = \frac{1}{20}, P_{101} = P_{011} = \frac{1}{10}, P_{003} = \frac{9}{20}$ , with all other  $P_{\vec{n}} = 0$ . The field at which the probabilities switch from one set of values to the other can be used to indicate when the energy levels change from the low to high field values. For  $^{40}\text{K}$  in a trap with 70kHz considered above, the shift  $l_r^3/v \approx -200$  corresponds to  $-\frac{1}{v} \approx 0.5 \times 10^{-7} a_0^{-3}$ , or a field  $\approx 1.15\text{G}$  above the free space resonance [8, 18], a large shift that should be readily discernible by experiments. This shift in field is roughly proportional to  $1/l_r^2$  hence  $\omega$ .

Finally, we discuss Feshbach resonances in higher angular momentum channels, say  $l, m_l$  (Bosons or non-identical Fermions if  $l$  even). Eq (5) is still valid, and the scattering amplitude has the form (generalizing eq (6))  $k^{2l} / \left( -\frac{1}{a_{lm}^{2l+1}} + c_{lm}k^2 + \dots - ik^{2l+1} \right)$ , where  $a_{lm}$  has dimension of a length. One have relations similar to eq (7) and (8) with coupling constants  $\alpha_{lm} = \frac{\sqrt{4\pi}}{(2l+1)!!} \int_{\vec{r}} \tilde{\varphi}_c(\vec{r}) r^l Y_{lm}^*(\hat{r})$ . We can obtain the corresponding part of  $G_E^{ref}$  with relevant  $l, m_l$  symmetry with  $g_l(r, r')$  given by an expression similar to eq(11) except that the  $b_1(r')$  ( $b_2(r')$ ) term has the factors  $\left(\frac{r}{l_r}\right)^l$  ( $\left(\frac{r}{l_r}\right)^{-(l+1)}$ ) with appropriate changes in the arguments of the confluent hypergeometric functions  $F$ , and  $h_1^{(1)} \rightarrow h_l^{(1)}$ . If we proceed as

in text and include only the regular term  $\propto r^l$  in  $g_l(r, r')$ , we would get

$$-\frac{1}{a_{lm}^{2l+1}} + c_{lm}k^2 + \dots = (-1)^{l+1} \left(\frac{2}{l_r}\right)^{2l+1} \frac{\Gamma(\frac{2l+3}{4} - \frac{E}{2\omega})}{\Gamma(-\frac{2l-1}{4} - \frac{E}{2\omega})} \quad (15)$$

This formula reduces to the known  $l = 0$  results in the literature [5, 6, 16] and the  $l = 1$  result above (eq (12)). However, this is incomplete if  $l \geq 2$ . For  $l = 2$  for example, the term  $\propto b_2$ , together with the  $in_2(kr)$  term from  $h_2^{(1)}(kr)$ , produce a term  $\propto \frac{1}{r^3} \times r^4 = r$  as the lowest order contribution to  $g_2(r, r')$ . (It can be verified that there are no terms of lower order in  $r$ ). As a result, the RHS of eq (15) acquires an extra term  $R_{2m}$  given by  $-\frac{2\Gamma(\frac{5}{2})}{l_r^4} \frac{\int_{\tilde{r}} \tilde{\varphi}_c(\tilde{r}) r Y_{2m}^*(\hat{r})}{\alpha_{2m}}$ . The integral involves short distance behavior of  $\tilde{\varphi}_c$ , and in contrast to the coupling constant  $\alpha_{lm}$ , cannot be eliminated from the final answer using the knowledge of the low energy scattering amplitude. Similar "residual" terms  $R_{lm}$  arise for all  $l \geq 2$ , though the expressions become more involved and will not be given here. However, fortunately, for a given  $lm$ ,  $R_{lm}$  is independent of the energy, magnetic field (if near resonance) and the energy level under consideration. Since near the resonance,  $\frac{1}{a_{lm}^{2l+1}}$  is expected to be linear in the deviation of  $B$  from resonance, the extra term  $R_{lm}$  simply gives a shift of the resonant field common to all the energy levels for given  $lm$  and trap parameters  $l_r$ . Apart from this, the behavior of the energy levels as a function of field is thus rather similar to that depicted in Fig 1 (though with energy levels given by  $(2n + \frac{2l+3}{2})\omega$  far from resonance). The quantitative details will of course depend on the values of the parameters  $c_{lm}$  etc.

In conclusion, we have calculated energy levels of two Fermions in a harmonic potential well under a finite angular momentum Feshbach resonance. In contrast to wide s-wave Feshbach resonances, there are large shifts in the resonant fields dependent on the energy levels. This prediction can be tested by experiments similar to those already performed in [3, 4].

This research is supported by the National Science Council of Taiwan under grant number NSC 95-2112-M-001-054-MY3.

- 
- [1] M. Greiner *et al*, Nature (London) **415**, 39 (2002)
  - [2] W. Hofstetter *et al*, Phys. Rev. Lett. **89**, 220407 (2002)
  - [3] M. Köhl *et al*, Phys. Rev. Lett. **94**, 080403 (2005)
  - [4] T. Stöferle *et al*, Phys. Rev. Lett. **96**, 030401 (2006)



- [5] D. B. M. Dickerscheid *et al*, Phys. Rev. A **71**, 043604 (2005)
- [6] R. B. Diener and T.-L. Ho, Phys. Rev. Lett. **96**, 010402 (2006)
- [7] L.-M. Duan, Phys. Rev. Lett. **95**, 243202 (2005)
- [8] C. Ticknor *et al*, Phys. Rev. A, **69**, 042712 (2004)
- [9] J. Zhang *et al*, Phys. Rev. A **70**, 030702(R) (2004)
- [10] J. P. Gaebler *et al*, Phys. Rev. Lett. **98**, 200403 (2007)
- [11] T. L. Ho and R. B. Diener, Phys. Rev. Lett. **94**, 090402 (2005)
- [12] V. Gurarie, L. Radzihovsky and A. V. Andreev, Phys. Rev. Lett. **94**, 230403 (2005)
- [13] C.H. Cheng and S.-K. Yip, Phys. Rev. Lett. **95**, 070404 (2005).
- [14] C.H. Cheng and S.-K. Yip, Phys. Rev. B, **73**, 064517 (2006).
- [15] D. Jaksch *et al*, Phys. Rev. Lett. **89**, 040402 (2002)
- [16] T. Busch *et al*, Found. Phys. **28**, 549 (1998)
- [17] H. G. Katzgraber *et al*, Phys. Rev. A **74**, 043602 (2006)
- [18] See also Fig 2 in ref [14]. The unit of the y-axis there should read  $10^{-7}a_0^{-3}$ . The correct values were however used in the rest of that paper.




FIG. 1: Energy levels versus  $-1/v$ . Energies are in units of the trap frequency,  $v$  is unit of  $l_r^3$ .




FIG. 2: The probabilities  $P$  of finding a particle in the state  $\vec{n}$  for  $\vec{n} = (000)$  (solid),  $(001)$  (dashed),  $(100)$  (long dashed), and  $(002)$  (dot-dashed), under the field sweep procedure described in text.  $v$  is unit of  $l_r^3$ .